

Temptra* (acetaminophen oral solution, USP)

Product Features

For fast, effective fever and pain relief.

<p>INFANT TEMPRA* Drops 80 mg/ 1 mL</p>	<p>CHILDREN'S TEMPRA* Syrup Regular Strength 80 mg/ 5 mL</p>	<p>Double-Strength CHILDREN'S TEMPRA* Syrup 160 mg/ 5 mL</p>
<ul style="list-style-type: none"> ▪ Alcohol, sugar and ASA free ▪ Cherry and dye-free banana flavours 	<ul style="list-style-type: none"> ▪ Alcohol and ASA free ▪ cherry flavour 	<ul style="list-style-type: none"> ▪ Alcohol and ASA free ▪ cherry and dye-free banana flavours

Dosage

Administer single dose orally, every 4-6 hours, according to the weights listed below with a daily maximum of 5 doses/24 hours.

Dosage ¹		INFANT TEMPRA* Drops 80 mg/ 1 mL	CHILDREN'S TEMPRA* Syrup Regular Strength 80 mg/ 5 mL	Double-Strength CHILDREN'S TEMPRA* Syrup 160 mg/ 5 mL
lbs	kg	0 – under 2 years Use supplied dropper only	2 - 6 years Use supplied dosage cup only	4 - 10 years Use supplied dosage cup only
6 - 11	2.0 - 5.4	Consult a physician	-	-
12 - 17	5.5 - 7.9	1.0 mL	-	-
18 - 23	8.0 – 10.9	1.5 mL	-	-
24 - 35	11.0 - 15.9	-	10 mL	5 mL
36 - 47	16.0 - 21.9	-	15 mL	7.5 mL
48 - 59	22.0 - 26.9	-	-	10 mL
60 - 71	27.0 - 31.9	-	-	12.5 mL
72 - 95	32.0 - 43.9	-	-	15 mL

¹These doses are based on weight, not age, and offer more specific doses than in the general instructions on label/carton.

WARNINGS

DO NOT USE with other drugs containing **acetaminophen**.

DO NOT take more than the recommended dose unless advised by a doctor. Taking more than the maximum daily dose may cause **severe or possibly fatal liver damage**.

Do not take if allergic to acetaminophen.

Consult a doctor if:

- you develop allergic reactions such as wheezing, rash or itching;
- your symptoms last for more than five days, or fever lasts more than 3 days.

OVERDOSE: In the case of an overdose, even if there are no signs or symptoms, call your Poison Control Centre or physician immediately. Within the first 24 hours your infant or child may experience nausea or vomiting, increased sweating, diarrhea, loss of appetite and stomach pain. Medical attention is critical, even in the absence of symptoms.

- **KEEP OUT OF THE REACH OF CHILDREN.** Safety sealed bottle. Do not use if seal is broken.

Bristol-Myers Squibb Canada
Montréal, Canada
Tel. No.

*Registered TM of E.R. Squibb & Sons LLC used under license by Bristol-Myers Squibb Canada

TMTM of E.R. Squibb & Sons LLC used under license by Bristol-Myers Squibb Canada

Date of Revision: 30 April 2010